

ELLIPTIC STARS IN A CHAOTIC NIGHT

T. Jäger*

March 29, 2010

Abstract

A recurrent theme in the description of phase portraits of dynamical systems is that of *elliptic islands in a chaotic sea*. Usually this picture is invoked in the context of smooth twist maps of the annulus or the torus, like the standard map. In this setting ‘elliptic islands’ refers to the topological disks bounded by periodic smooth curves surrounding elliptic periodic points. Establishing the existence of these curves is one of the many achievements of KAM-theory.

The aim of this note is to approach the topic from a different angle, namely from the viewpoint of rotation theory in a purely topological setting. We study homeomorphisms of the two-torus, homotopic to the identity, which have no wandering open sets (as in the area-preserving case) and whose rotation set has non-empty interior. We define local rotation subsets $\rho_F(U)$ by restricting Misiurewicz and Ziemian’s definition of the rotation set to starting points in a small open disk U . Our main result is the following dichotomy: Either $\rho_U(F)$ is reduced to a single rational vector and U is contained in a periodic topological open disk which contains a periodic point, or $\rho_U(F)$ is large, in the sense that its convex hull has non-empty interior. This allows to distinguish an ‘elliptic’ and a ‘chaotic’ regime, and as a consequence we obtain that in the chaotic region the dynamics are sensitive with respect to initial conditions.

In order to demonstrate these results we introduce a parameter family of smooth toral diffeomorphisms that is inspired by an example of Misiurewicz and Ziemian. The pictures obtained from simulations in this family motivate an alternative formulation of the original theme.




Figure 1: Elliptic islands surrounding two 2-periodic orbits of the map $f_\alpha(x, y) = (x + \alpha \sin(2\pi(y + \alpha \sin(2\pi x))), y + \alpha \sin(2\pi x))$ with $\alpha = 0.5$.

*TU Dresden, Institut für Analysis. Email: Tobias.Oertel-Jaeger@tu-dresden.de

1 Introduction

We denote by $\text{Homeo}_0(\mathbb{T}^2)$ the set of homeomorphisms of the two-torus that are homotopic to the identity. Given a lift $F : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ of $f \in \text{Homeo}_0(\mathbb{T}^2)$, Misiurewicz and Ziemian [1] introduced the rotation set of F as

$$(1.1) \quad \rho(F) := \left\{ \rho \in \mathbb{R}^2 \mid \exists n_i \nearrow \infty, z_i \in \mathbb{R}^2 : \lim_{i \rightarrow \infty} (F^{n_i}(z_i) - z_i) / n_i = \rho \right\}.$$

This set is always compact and convex [1]. Further, the properties of $\rho(F)$ have strong implications for the dynamics of f . In particular, this is true for the situation we will concentrate, namely when $\rho(F)$ has non-empty interior. In this case all rotation vectors in $\text{int}(\rho(F))$ are realised on minimal sets [2, 3, 4] and the topological entropy of f is strictly positive [5]. Further the set $\{f \in \text{Homeo}_0(\mathbb{T}^2) \mid \text{int}(F) \neq \emptyset\}$ is an open and therefore, in a topological sense, large subset of $\text{Homeo}_0(\mathbb{T}^2)$ [3].

Our aim is to give some meaning to the notion of *elliptic islands in a chaotic sea* in this purely topological setting. To that end, we restrict the definition in (1.1) to orbits starting in some subset $U \subseteq \mathbb{T}^2$. Let $\pi : \mathbb{R}^2 \rightarrow \mathbb{T}^2$ denote the canonical projection. We define the *rotation subset on U* by

$$(1.2) \quad \rho_U(F) := \left\{ \rho \in \mathbb{R}^2 \mid \exists z_i \in \pi^{-1}(U), n_i \nearrow \infty : \lim_{i \rightarrow \infty} (F^{n_i}(z_i) - z_i) / n_i = \rho \right\}.$$

In general, even when U is open $\rho_U(F)$ can be much smaller than $\rho(F)$. For instance, when f is a sufficiently smooth toral diffeomorphism then generic elliptic periodic points are surrounded by periodic invariant curves (see, for example, [6, 9]). The rotation subsets of the corresponding topological disks contain a single rational rotation vector, whereas $\rho(F)$ may have non-empty interior. A more general example is sketched in Remark 1.2 below.

However, when U is open and recurrent, then in a number of situations $\rho(F)$ is already determined by $\rho_U(F)$. In order to give precise statements, we need some notation and terminology. We say $U \subseteq \mathbb{T}^2$ is *bounded* if the connected components of its lift to $\mathbb{R}^2$ are bounded. Given $f \in \text{Homeo}_0(\mathbb{T}^2)$ we say U is *wandering* if $f^n(U) \cap U = \emptyset \forall n \geq 1$ and *non-wandering* otherwise. We call U *recurrent* if there exist infinitely many $n \in \mathbb{N}$ with $f^n(U) \cap U \neq \emptyset$. We call $z \in \mathbb{T}^2$ *wandering* if it is contained in some wandering open set and *non-wandering* otherwise. It is easy to see that if U is open and contains a non-wandering point then it is recurrent. Finally, we say that f is *non-wandering* if it has all points non-wandering. In this case all open sets are recurrent. Note that any area-preserving toral homeomorphism is non-wandering.

Given a lift F of $f \in \text{Homeo}_0(\mathbb{T}^2)$, let $\varphi_n(z) = (F^n(z) - z) / n$. If $\lambda \in \mathbb{R}$ and $v \in \mathbb{R}^2 \setminus \{0\}$, let $L_{\lambda,v} = \lambda v + \{v\}^\perp$.

Theorem 1.1. *Suppose F is a lift of $f \in \text{Homeo}_0(\mathbb{T}^2)$ and $U \subseteq \mathbb{T}^2$ is open, bounded, connected and recurrent.*

- (a) *If $\rho_U(F) \subseteq L_{\lambda,v}$ for some $\lambda \in \mathbb{R}$, $v \in \mathbb{R}^2 \setminus \{0\}$, then either $\rho(F) \subseteq L_{\lambda,v}$ or $\rho_U(F)$ is reduced to a single rational vector.*
- (b) *If $\mathcal{S}$ is a line segment of positive length without rational points and $\rho_U(F) = \mathcal{S}$, then $\rho(F) = \mathcal{S}$. Further, $\varphi_n(U)$ converges to $\mathcal{S}$ in Hausdorff distance as $n \rightarrow \infty$.*
- (c) *If $\rho_U(F) = \{\rho\}$ with $\rho \in \mathbb{R}^2$ irrational¹ then $\rho(F) = \{\rho\}$.*

Remark 1.2. We note that when no recurrence assumption is made no relation between $\rho_U(F)$ and $\rho(F)$ can be expected. Without going into detail, we want to mention a possible way to construct respective examples: When $\rho(F)$ has non-empty interior, then for any compact connected subset $C \subseteq \rho(F)$ there exists a point $z \in \mathbb{T}^2$ with $\rho_{\{z\}}(F) = C$ [5]. By blowing up the points in the orbit of z to small disks in a Denjoy-like construction one may thus obtain a wandering open set U whose rotation set is an arbitrary compact connected subset of the rotation set.

Theorem 1.1(i) implies that if f is non-wandering and $\rho(F)$ has non-empty interior then the rotation subset of U can only be contained in a line if it is reduced to a single rational rotation vector, that is, $\rho_U(F) = \{\rho\}$ with $\rho \in \mathbb{Q}^2$. Together with some additional details on the rational case, this yields our main result.

Theorem 1.3. *Let $F : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a lift of $f \in \text{Homeo}_0(\mathbb{T}^2)$ and suppose that f is non-wandering and $\rho(F)$ has non-empty interior. Then for any open, bounded and connected set U one of the following two holds.*

- (i) *$\rho_U(F)$ is reduced to a single rational vector ρ and U is contained in an embedded topological open disk $D \subseteq \mathbb{T}^2$ which is invariant under some iterate f^p and contains a p -periodic point.*
- (ii) *The convex hull of $\rho_U(F)$ has non-empty interior.*

¹We call a vector $\rho = (\rho_1, \rho_2) \in \mathbb{R}^2$ *rational* if $\rho \in \mathbb{Q}^2$, *irrational* if $\rho_1, \rho_2, \rho_1/\rho_2 \notin \mathbb{Q}$ and *semi-rational* if it is neither rational nor irrational.

The above result allows to give an intrinsic definition of ‘elliptic’ and ‘chaotic’ regions. Given a set $A \subseteq \mathbb{R}^2$ we denote by $\text{Conv}(A)$ its convex hull and by $\text{int}(A)$ its interior.

Definition 1.4. Suppose $f \in \text{Homeo}_0(\mathbb{T}^2)$. Let

$$\begin{aligned}\mathcal{E}(f) &:= \{z \in \mathbb{T}^2 \mid \# \rho_U(F) = 1 \text{ for some open neighbourhood } U \text{ of } z\} , \\ \mathcal{C}(f) &:= \{z \in \mathbb{T}^2 \mid \text{int}(\text{Conv}(\rho_U(F))) \neq \emptyset \text{ } \forall \text{ open neighbourhoods } U \text{ of } z\} .\end{aligned}$$

A point $z \in \mathbb{T}^2$ is called ε -Lyapunov stable if there exists some $\delta > 0$ such that $f^n(B_\delta(z)) \subseteq B_\varepsilon(f^n(z)) \forall n \in \mathbb{N}$ and Lyapunov stable if it is ε -Lyapunov stable for all $\varepsilon > 0$. As one should expect for a notion of stability, Lyapunov stable points do not occur in the ‘chaotic’ regime.

Proposition 1.5. Suppose $f \in \text{Homeo}_0(\mathbb{T}^2)$ and $\rho(F)$ has non-empty interior.

- (a) If f is non-wandering then no point in $\mathcal{C}(f)$ is $\frac{1}{2}$ -Lyapunov stable.
- (b) If f is area-preserving, U is a connected and bounded neighbourhood of $z \in \mathcal{C}(f)$ and $\hat{U}$ is a connected component of $\pi^{-1}(U)$ then $\limsup_{n \rightarrow \infty} \text{diam} \frac{1}{n} (F^n(\hat{U})) > 0$.

Note that in contrast to this, in the construction sketched in Remark 1.2 all points in the wandering topological disks will be Lyapunov stable provided the diameter of these disks goes to zero along the orbit.

The paper is organised as follows: In Section 2 we collect a number of basic statements on rotation subsets. Section 3 then contains the technical core of the paper. We work on the universal cover $\mathbb{R}^2$ and consider bounded open and connected sets that intersect their image. In this setting, we describe a number of situations in which the rotation subset already determines the rotation set, or at least forces it to be contained in a line segment. In Section 4 these statements are then used to prove the main results. Finally, in Section 5 we provide some explicit examples to which our results apply. To that end, we introduce a parameter family of smooth torus diffeomorphisms that is based on an example by Misiurewicz and Ziemian in [3]. For appropriate parameter values these maps have a rotation set with non-empty interior, and simulations clearly indicate that they also exhibit elliptic islands.

Acknowledgements. This work was supported by an Emmy Noether Grant (Ja 1721/2-1) of the German Research Council (DFG). The results were partially presented at the Workshop on Dynamics in Dimension two, April 2009 in Pucón (Chile). I would like to thank the organisers Mario Ponce and Andres Navas for the great opportunity they created. I am further indebted to Andres Koropecki and Patrice Le Calvez for thoughtful remarks and stimulating discussions and to Hendrik Vogt for drawing my attention to the fine structure of the elliptic islands depicted in Figure 6. Finally, I would like to thank Jean-Christophe Yoccoz for his support during my time at the Collège de France, where a part of this work was carried out.

2 Some basic results on rotation subsets

The aim of this section is to collect a number of elementary statements on rotation subsets and rotation vectors that will be used in the later sections. For the purposes of this section there is no need to restrict to dimension 2. Hence, we will work on $\mathbb{T}^d$ ($d \in \mathbb{N}$), with the definitions of the rotation set and rotation subsets analogous to those on $\mathbb{T}^2$.

Notation. We denote the Euclidean scalar product of vectors $v, w \in \mathbb{R}^d$ by $\langle v, w \rangle$ and also write $\langle w \rangle_v$ instead of $\langle v, w \rangle$. By $\|v\| = \sqrt{\langle v, v \rangle}$ we denote the Euclidean length of the vector v . If G is an additive group then $G_* = G \setminus \{0\}$. By $\text{Conv}(C)$ we denote the convex hull of a subset $C \subseteq \mathbb{R}^d$. By $\text{Ex}(C)$ we denote the extremal points of $\text{Conv}(C)$ and let $\text{Conv}_\times(C) = \text{Conv}(C) \setminus \text{Ex}(C)$.

If $v = (v_1, v_2) \in \mathbb{R}^2$ we let $v^\perp = (-v_2, v_1)$. Given $\lambda \neq 0$, $v \in \mathbb{R}_*^2$ and $a \leq b \in \mathbb{R} \cup \{\pm\infty\}$ we let $L_{\lambda,v} = \lambda v + \{v\}^\perp$ and

$$\begin{aligned}C_v[a, b] &= \{z \in \mathbb{R}^2 \mid a\langle z, v \rangle \leq \langle z, v^\perp \rangle \leq b\langle z, v \rangle\} \\ L_{\lambda,v}[a, b] &= \{z \in L_{\lambda,v} \mid a\langle z, v \rangle \leq \langle z, v^\perp \rangle \leq b\langle z, v \rangle\} .\end{aligned}$$

Note that thus $L_{\lambda,v}(a, b) = L_{\lambda,v} \cap C_v(a, b)$ and $C_v[a, a] = \mathbb{R} \cdot (v + av^\perp)$. Further, we let

$$\begin{aligned}C_v^+[a, b] &= \{z \in C_v[a, b] \mid \langle z, v \rangle \geq 0\} \text{ and} \\ S_v[a, b] &= \{z \in \mathbb{R}^2 \mid \langle z, v \rangle \in [a, b]\} .\end{aligned}$$

All these notions are used similarly for open and half-open intervals.

The following basic observation is a direct consequence of the definition in (1.2). Recall that $\varphi_n(z) = (F^n(z) - z)/n$.

Lemma 2.1. *Suppose F is a lift of $f \in \text{Homeo}_0(\mathbb{T}^d)$ and $U \subseteq \mathbb{T}^d$. Then for all $\varepsilon > 0$ there exists some $n_0 = n_0(\varepsilon) \in \mathbb{N}$ such that $\varphi_n(U) \subseteq B_\varepsilon(\rho_U(F)) \forall n \geq n_0$.*

The proof of the following statement is more or less identical to that of the connectedness of the rotation set in [1], but we include it for the convenience of the reader.

Lemma 2.2. *Let F be a lift of $f \in \text{Homeo}_0(\mathbb{T}^d)$. For any $U \subseteq \mathbb{T}^d$, the set $\rho_U(F)$ is compact. Further, if U is connected then so is $\rho_U(F)$.*

Proof of Lemma 2.2. The fact that $\rho_U(F)$ is compact follows immediately from the definition. Suppose for a contradiction that U is connected but $\rho_U(F)$ is not. Then there exist disjoint open sets V_1 and V_2 with $\rho_U(F) \subseteq V_1 \cup V_2$ and $\rho_U(F) \cap V_i \neq \emptyset$ ($i = 1, 2$). Since $\rho_U(F)$ is compact, we may assume that $\varepsilon = d(V_1, V_2) > 0$. Lemma 2.1 implies that there exists $n_0 \in \mathbb{N}$ such that $\varphi_n(U) \subseteq V_1 \cup V_2 \forall n \geq n_0$. Further, since $\sup_{z \in \mathbb{R}^2} \|F(z) - z\| < \infty$ there exists $n_1 \in \mathbb{N}$ such that

$$(2.1) \quad |(F^{n+1}(z) - z)/(n+1) - (F^n(z) - z)/n| < \varepsilon \quad \forall n \geq n_1, z \in \mathbb{R}^2.$$

It follows that if $n \geq n_1$ and $\varphi_n(U) \subseteq V_i$, then $\varphi_k(U) \subseteq V_i \forall k \geq n$ and therefore $\rho_U(F) \subseteq \overline{V_i}$. Since this is not the case, $\varphi_n(U)$ must intersect both V_1 and V_2 for all $n \geq n_1$. However, for $n \geq \max\{n_0, n_1\}$ we then obtain $\varphi_n(U) \subseteq V_1 \cup V_2$ and $\varphi_n(U) \cap V_i \neq \emptyset$ ($i = 1, 2$). This contradicts the connectedness of $\varphi_n(U)$. $\square$

Deviations from a constant rotation and invariant measures. For $f \in \text{Homeo}_0(\mathbb{T}^d)$ with lift $F : \mathbb{R}^d \rightarrow \mathbb{R}^d$, $\rho \in \mathbb{R}^d$ and $v \in \mathbb{R}_*^d$ we let

$$(2.2) \quad D_n(z, \rho) := F^n(z) - z - n\rho \quad \text{and} \quad D_n^v(z, \rho) := \langle D_n(z, \rho) \rangle_v.$$

If we need to make the dependence on f explicit, we also write $D_{f,n}(z, \rho)$ and $D_{f,n}^v(z, \rho)$. For any f -invariant probability measure μ , the rotation number of f with respect to μ is given by

$$(2.3) \quad \rho_\mu(F) = \int_{\mathbb{T}^d} F(z) - z \, d\mu(z).$$

When F is fixed and no ambiguities can arise, we suppress it from the notation and write ρ_μ instead of $\rho_\mu(F)$. By $\text{supp}(\mu)$ we denote the topological support of μ .

Lemma 2.3. *Suppose $F : \mathbb{R}^d \rightarrow \mathbb{R}^d$ is a lift of $f \in \text{Homeo}_0(\mathbb{T}^d)$ and μ is an ergodic f -invariant probability measure. Then there exists no constant $s > 0$ with the property that for μ -a.e. $z \in \mathbb{R}^2$ there is a positive integer n_z such that $D_{n_z}^v(z, \rho_\mu) \geq sn_z$.*

Proof. We suppose for a contradiction that a constant $s > 0$ with the above property exists. We fix an f -invariant set $\Omega \subseteq \mathbb{T}^2$ of measure $\mu(\Omega) = 1$ such that for all $z \in \Omega$ there exists $n_z \in \mathbb{N}$ with

$$(2.4) \quad D_{n_z}^v(z, \rho_\mu) \geq sn_z.$$

In addition, we assume that

$$(2.5) \quad \lim_{n \rightarrow \infty} (F^n(z) - z)/n = \rho_\mu \quad \forall z \in \Omega.$$

Given any $z_0 \in \Omega$, we recursively define a sequence of integers n_i by $n_0 = 0$ and $n_{i+1} = n_i + n_{F^{n_i}(z_0)}$. Then we obtain

$$\begin{aligned} D_{n_k}^v(z_0, \rho_\mu) &= \langle F^{n_k}(z_0) - z_0 - n_k \rho_\mu \rangle_v \\ &= \left\langle \sum_{i=0}^{k-1} F^{n_{i+1}}(z_0) - F^{n_i}(z_0) - (n_{i+1} - n_i) \rho_\mu \right\rangle_v \\ &= \left\langle \sum_{i=0}^{k-1} F^{n_{F^{n_i}(z_0)}}(F^{n_i}(z_0)) - F^{n_i}(z_0) - (n_{F^{n_i}(z_0)}) \rho_\mu \right\rangle_v \\ (2.4) \quad &\geq \sum_{i=0}^{k-1} s(n_{i+1} - n_i) = sn_k. \end{aligned}$$

Hence $\lim_{k \rightarrow \infty} D_{n_k}^v(z_0, \rho_\mu)/n_k \geq s$, contradicting (2.5) which implies $\lim_{n \rightarrow \infty} D_n^v(z_0, \rho_\mu) = 0$. $\square$

A reduction lemma. Any integer matrix $M \in \text{GL}(d \times d, \mathbb{Z})$ induces a toral endomorphism $g_M : \mathbb{T}^d \rightarrow \mathbb{T}^d$, $\pi(z) \mapsto \pi(Mz)$, and g_M is invertible if and only if $M \in \text{SL}(d, \mathbb{Z})$. The following lemma describes how a coordinate transformation by such a map g_M acts on the rotation set.

Lemma 2.4 (Reduction Lemma). *Suppose $f \in \text{Homeo}_0(\mathbb{T}^d)$ has lift $F : \mathbb{R}^d \rightarrow \mathbb{R}^d$, $U \subseteq \mathbb{T}^d$ and $M \in \text{SL}(d, \mathbb{Z})$. Then the following hold.*

(a) *Let $\tilde{f} = g_M^{-1} \circ f \circ g_M \in \text{Homeo}_0(\mathbb{T}^d)$ with lift $\tilde{F} = M^{-1} \circ F \circ M$. Then*

$$(2.6) \quad \rho_{g_M^{-1}U}(\tilde{F}) = M^{-1}(\rho_U(F)) .$$

(b) *If $\rho_U(F) \subseteq L_{\lambda,v}$ then $\rho_{g_M^{-1}U}(\tilde{F}) \subseteq M^{-1}(L_{\lambda,v}) = L_{\tilde{\lambda},\tilde{v}}$, where $\tilde{v} = M^t v$ and $\tilde{\lambda} = \lambda \|v\|^2 / \|\tilde{v}\|^2$. Further $D_{f,n}^v(z, \rho) = D_{\tilde{f},n}^{\tilde{v}}(M^{-1}z, M^{-1}\rho) \forall n \in \mathbb{N}, z, \rho \in \mathbb{R}^d$.*

(c) *Let $1 \leq k < d$ and suppose that $w_1, \dots, w_k$ are linearly independent integer vectors and $\text{Conv}_\times(\{w_1, \dots, w_k\})$ contains no further integer vectors. (If $k = 1$ this just means that the entries of w_1 are relatively prime.) Then there exist integer vectors $w_{k+1}, \dots, w_d$ such that $\det(w_1, \dots, w_d) = 1$.*

Note that for any integer vector $w \in \mathbb{Z}^d$ with relatively prime entries part (c) allows to perform a linear coordinate transformation on $\mathbb{T}^d$ such that w becomes a base vector.

Proof.

(a) Suppose $z_i \in \pi^{-1}U$, $n_i \nearrow \infty$ and $\lim_{i \rightarrow \infty} (F^{n_i}(z_i) - z_i) / n_i = \rho$. Then

$$\begin{aligned} M^{-1}\rho &= M^{-1} \left(\lim_{i \rightarrow \infty} (F^{n_i}(z_i) - z_i) / n_i \right) \\ &= \lim_{i \rightarrow \infty} (M^{-1} \circ F^{n_i} \circ M (M^{-1}z_i) - M^{-1}z_i) / n_i \\ &= \lim_{i \rightarrow \infty} (\tilde{F}^{n_i}(M^{-1}z_i) - M^{-1}z_i) / n_i \in \rho_{M^{-1}U}(\tilde{F}) . \end{aligned}$$

This shows that $M^{-1}(\rho_U(F)) \subseteq \rho_{g_M^{-1}U}(\tilde{F})$ and since $f = g_M \circ \tilde{f} \circ g_M^{-1}$ the opposite inclusion follows in the same way.

(b) $\rho_{g_M^{-1}U}(\tilde{F}) \subseteq M^{-1}(L_{\lambda,v})$ holds by part (a). Further, we have $M^{-1}(L_{\lambda,v}) = M^{-1}(\lambda v) + M^{-1}\{v\}^\perp$. Since $\tilde{v} = M^t v \perp M^{-1}\{v\}^\perp$ it follows that $M^{-1}(L_{\lambda,v}) = L_{\tilde{\lambda},\tilde{v}}$ for some $\tilde{\lambda} \in \mathbb{R}$ and we have $\tilde{\lambda} \|\tilde{v}\| = \langle M^{-1}(\lambda v), \tilde{v} / \|\tilde{v}\| \rangle = (\lambda / \|\tilde{v}\|) \cdot \langle v, v \rangle$. Finally, in order to check that $D_{f,n}^v(z, \rho) = D_{\tilde{f},n}^{\tilde{v}}(M^{-1}z, M^{-1}\rho)$ let $z \in \mathbb{R}^d$. Then

$$\begin{aligned} D_{\tilde{f},n}^{\tilde{v}}(M^{-1}z, M^{-1}\rho) &= \langle \tilde{F}^n(M^{-1}z) - M^{-1}z - nM^{-1}\rho, \tilde{v} \rangle \\ &= \langle M^{-1} \circ F^n(z) - M^{-1}z - nM^{-1}\rho, M^t v \rangle \\ &= \langle F^n(z) - z - n\rho, v \rangle = D_{f,n}^v(z, \rho) . \end{aligned}$$

(c) Choose integer vectors $w_{k+1}, \dots, w_d$ such that $\text{span}(w_1, \dots, w_d) = \mathbb{R}^d$. Then $|\det(w_1, \dots, w_d)| \geq 2$ if and only if $\text{Conv}_\times(w_1, \dots, w_d)$ contains an integer vector. In this case we replace one of the vectors $w_{k+1}, \dots, w_d$ by an integer vector in $\text{Conv}_\times(w_1, \dots, w_d)$ such that the new set of vectors still spans $\mathbb{R}^d$. This reduces the absolute value of the determinant, and after a finite number of steps we arrive at $\det(w_1, \dots, w_d) = \pm 1$. Replacing w_d by $-w_d$ if necessary we obtain $\det(w_1, \dots, w_d) = 1$. $\square$

3 Rotation subsets on the universal cover

Throughout this section, we suppose that G is the lift of a toral homeomorphism $g \in \text{Homeo}_0(\mathbb{T}^2)$, $\hat{U} \subseteq \mathbb{R}^2$ is bounded and connected and $G(\hat{U}) \cap \hat{U} \neq \emptyset$. Further, we assume that $\lambda \neq 0$ and $v \in \mathbb{R}_*^2$.

In order to control the whole rotation set by using assumptions on $\rho_{\hat{U}}(G)$, we proceed in several steps. The first is to obtain some information about the extremal points of the rotation set.

Lemma 3.1. *Suppose $\rho_{\hat{U}}(G) = L_{\lambda,v}[a, b]$. Then all extremal points of $\rho(G)$ belong to $C_v[a, b]$.*

Proof. Suppose that $\lambda > 0$. (Otherwise we replace v by $-v$.) Performing a linear change of coordinates via Lemma 2.4 if necessary we may assume that

$$(3.1) \quad \pi_1(L_{\lambda,v}[a, b]) \subseteq (0, \infty) ,$$

such that in particular $C_v^+[a, b] \setminus \{0\} \subseteq (0, \infty) \times \mathbb{R}$. Further, we may assume that $\widehat{U}$ intersects $\{0\} \times \mathbb{R}$, otherwise we replace it by an integer translate and/or one of its iterates.² Let $V := \bigcup_{n \in \mathbb{N}_0} G^n(\widehat{U})$. As $G(\widehat{U}) \cap \widehat{U} \neq \emptyset$, the set V is connected. We claim that for sufficiently large $l \in \mathbb{N}$ the integer translate $V - (0, l)$ is disjoint from V . In order to see this let $r := \sup_{z \in \widehat{U}} \|z\|$. Due to (3.1) and Lemma 2.1 only a finite number of iterates of $\widehat{U}$ intersect $[-r, r] \times \mathbb{R}$. Therefore V and $\widehat{U} - (0, l)$ are disjoint for large l . Hence, if the orbit of $\widehat{U} - (0, l)$ intersects V then it must first intersect $\widehat{U}$. However, by the same argument the orbit of $\widehat{U} - (0, l)$ can only intersect a finite number of its vertical integer translates, such that for sufficiently large l we have $V \cap [V - (0, l)] = \emptyset$ as required.

Now let $y_1 = \inf\{y \in \mathbb{R} \mid (0, y) \in V - (0, l)\}$, $y_2 = \sup\{y \in \mathbb{R} \mid (0, y) \in V\}$ and define W as the union of V , $V - (0, l)$ and the vertical arc from $(0, y_1)$ to $(0, y_2)$. Let Y be the unique connected component of $\mathbb{R}^2 \setminus \overline{W}$ which is unbounded to the left and $A = \mathbb{R}^2 \setminus Y$.




Figure 2: Construction of the sets W on the left and A on the right.

The following three remarks about these objects will be helpful. First, as $\rho_{\widehat{U}}(G) \subseteq L_{\lambda, v}[a, b]$ and $\lambda > 0$, we have that for all $\alpha < a$ and $\beta > b$ there exists a constant $R = R(\alpha, \beta) > 0$ such that

$$(3.2) \quad A \subseteq B_R(C_v^+(\alpha, \beta)) .$$

Secondly, due to the definition of W , its connectedness and (3.1), the set $Z := (\mathbb{R}^+ \times \mathbb{R}) \setminus A = (\mathbb{R}^+ \times \mathbb{R}) \cap Y$ consists of exactly two connected components. These can be defined as follows. Fix any $\zeta_0 \in Y$ with $\pi_1(\zeta_0) < 0$. For any $\zeta \in Z$, there is a path γ_ζ in Y from ζ to ζ_0 . Let y_ζ be the second coordinate of the first point in which γ_ζ intersects the vertical axis. The fact whether y_ζ lies below y_1 or above y_2 does not depend on the choice of the path, since this would contradict the connectedness of W . Hence, $Z^- = \{\zeta \in Z \mid y_\zeta < y_1\}$ and $Z^+ = \{\zeta \in Z \mid y_\zeta > y_2\}$ form a partition of Z into two connected components.

Thirdly, there holds $\mathbb{R}^+ \times \mathbb{R} \subseteq \bigcup_{k \in \mathbb{N}} A + (0, kl)$. Consequently, for any $m \in \mathbb{N}$ the set $A \cap \pi_1^{-1}[m, m+1)$ contains a fundamental domain of $\mathbb{T}^2$, that is, $\pi(A \cap \pi_1^{-1}[m, m+1)) = \mathbb{T}^2$.

It is important to note, however, that A is not G -invariant. Yet, in order to obtain control over the full rotation set via A we will need to ensure that orbits ‘moving to the right’ become ‘trapped’ in A (or one of its integer translates). Hence, the following statement is crucial for our purposes.

Claim 3.2. *There exists a constant $K > 0$ such that $z \in A \cap \pi_1^{-1}[K, \infty)$ implies $G^{\pm 1}(z) \in A$.*

Proof. We show that there exists a constant $K' > 0$ such that $z \in Z \cap \pi_1^{-1}[K', \infty)$ implies $G^{\pm 1}(z) \in Z$. If we let

$$(3.3) \quad M := \sup_{z \in \mathbb{R}^2} \|G(z) - z\| = \sup_{n \in \mathbb{R}^2} \|G^{-1}(z) - z\| ,$$

then for any $z \in \pi_1^{-1}[K' + M, \infty)$ this means that $G^{\pm 1}(z) \in Z$ implies $z \in Z$ and hence $z \in A$ implies $G^{\pm 1}(z) \in A$. Thus we can choose $K = K' + M$.

²Note that by assumption $\bigcup_{n \in \mathbb{N}_0} G^n(\widehat{U})$ is connected and $\pi_1 \circ G^n(\widehat{U})$ goes to ∞ as $n \rightarrow \infty$ due to (3.1). Hence, one of the iterates of $\widehat{U}$ has to intersect an integer vertical $\{m\} \times \mathbb{R}$.

From (3.1) and Lemma 2.1 we deduce that there exists $n_0 \in \mathbb{N}$ such that $\pi_1 \circ G^n(\widehat{U}) \subseteq (4M, \infty) \forall n \geq n_0$. Let $K' > 0$ such that for all $j \leq n_0$ there holds $\pi_1 \circ G^j(\widehat{U}) \subseteq [0, K']$. Then we have

$$(3.4) \quad \pi_1 \circ G^n(\widehat{U}) \cap [K', \infty) \neq \emptyset \quad \Rightarrow \quad \pi_1 \circ G^k(\widehat{U}) \cap [0, 4M] = \emptyset \quad \forall k \geq n.$$

The same statement applies to $\widehat{U} + (0, l)$. Due to (3.2) there exists $C > 0$ such that

$$B := [0, 4M] \times [C - M, \infty) \subseteq Z^+.$$

Let $z^* = (3M, C)$ and fix $z \in Z^+$ with $\pi_1(z) \geq K'$. Then, since Z^+ is open and connected, there is a simple path $\gamma : [0, 1] \rightarrow Z^+$ from z to z^* . We claim that γ can be chosen such that its image is contained in $Z^+ \cap \pi_1^{-1}[3M, \infty)$. Suppose not and let $t_0 := \min\{t \in [0, 1] \mid \gamma(t) \in B\}$ and $\Gamma = \{\gamma(t) \mid t \in [0, t_0]\}$. Then Γ divides the set $([0, \pi_1(z)] \times \mathbb{R}) \setminus B$ into exactly two connected components D^+ and D^- that are unbounded above, respectively below. Now, if $\overline{W}$ does not




Figure 3: The domains D^- and D^+ .

intersect $D^+ \cap \pi_1^{-1}[0, 4M)$ then $D^+ \cap \pi_1^{-1}[0, 4M) \subseteq Z^+$, and it is easy to see that in this case either γ does not intersect $\pi_1^{-1}[0, 3M)$ or we can modify it to that end. Otherwise, there must be some $k \in \mathbb{N}$ such that $G^k(\widehat{U})$ or $G^k(\widehat{U}) - (0, l)$ intersects $D^+ \cap \pi_1^{-1}[0, 4M)$. However, since $\widehat{U}$ intersects D^- and the set $\bigcup_{i=0}^k G^i(\widehat{U})$ is connected, this implies that there must be some $n \leq k$ such that $\pi_1 \circ G^n(\widehat{U})$ intersects $[\pi_1(z), \infty) \subseteq [K', \infty)$. This contradicts (3.4).

Summarising, we have found a path γ from z to z^* which is contained in $Z^+ \cap \pi_1^{-1}[3M, \infty)$. In particular, γ is contained in the complement of $\overline{W}$. Consequently, the path $G \circ \gamma$ is contained in the complement of $G(\overline{W})$. At the same time, it is also contained in $\pi_1^{-1}[2M, \infty)$. However, it follows from the construction of W and the definition of M that $G(\overline{W}) \cap \pi_1^{-1}[2M, \infty) = \overline{W} \cap \pi_1^{-1}[2M, \infty)$. Hence, the path $G \circ \gamma$ is contained in the complement of $\overline{W}$ as well. Furthermore, it joins $G(z)$ to the point $G(z^*)$. Since the latter is contained in $B \subseteq Z^+$, this implies that $G(z)$ is equally contained in Z^+ . When $z \in Z^-$ the argument is similar. In the same way one can show that $G^{-1}(z) \in Z$, and this proves the claim. $\circ$

In order to complete the proof of Lemma 3.1, suppose that ρ is an extremal point of $\rho(G)$ which is not contained in $C_v[a, b]$. By performing a linear change of coordinates again if necessary, we may assume that $\pi_1(\rho) > 0$.³ Since ρ is realised by an ergodic invariant measure [1, Corollary 3.5], there exists a point $z_0 \in \mathbb{R}^2$ with

$$(3.5) \quad \lim_{n \rightarrow \infty} (G^n(z_0) - z_0)/n = \rho.$$

³Choose a basis of integer vectors w_1, w_2 with $\det(w_1, w_2) = 1$ such that both w_1 and ρ lie to the right of the oriented line $\mathbb{R}w_2$ and $w_2 \notin C_v[a, b]$ (the latter ensures that (3.1) remains valid.) Then apply Lemma 2.4.

Let $z_n = G^n(z_0)$ and $\mathcal{O}^+(z_0) = \{z_n \mid n \geq 0\}$. Then, due to (3.5), for every $\gamma < 0 < \delta$ there exists $\tilde{R} > 0$ such that $\mathcal{O}^+(z_0) \subseteq z_0 + B_{\tilde{R}}(C_v^+[\gamma, \delta])$. If γ, δ are chosen sufficiently close to 0 and α, β in (3.2) are sufficiently close to a and b then $B_R(C_v[\alpha, \beta]) \cap (z_0 + B_{\tilde{R}}(C_v^+[\gamma, \delta]))$ is bounded (recall that ρ is not contained in $C_v[a, b]$). Further, by replacing z_0 with an integer translate if necessary, we may assume that $z_0 \in A$ and $\pi_1(z_0) \geq K + \tilde{R}$, where K is chosen as in Claim 3.2. It follows that $\pi_1(z_n) \geq K$ and hence $z_n \in A \forall n \in \mathbb{N}$. Consequently $\mathcal{O}^+(z_0) \subseteq B_R(C_v[\alpha, \beta]) \cap (z_0 + B_{\tilde{R}}(C_v^+[\gamma, \delta]))$ such that $\mathcal{O}^+(z_0)$ is bounded, contradicting (3.5). Hence, all extremal points of $\rho(G)$ must be contained in $C_v[a, b]$. $\square$

In the opposite way, information about the extremal points of $\rho(G)$ allows to draw conclusions about the behaviour of the iterates of $\hat{U}$.

Lemma 3.3. *Suppose that $\rho_{\hat{U}}(G) = L_{\lambda, v}[a, b]$, $\gamma \in [a, b]$ and $\rho \in C_v[\gamma, \gamma]$ is an extremal point of $\rho(G)$. Then given any $\varepsilon > 0$ there exists $N \in \mathbb{N}$ such that $G^n(\hat{U}) \cap C_v^+(\gamma - \varepsilon, \gamma + \varepsilon) \neq \emptyset \forall n \geq N$.*

Proof. As in the proof of Lemma 3.1 we assume $\lambda > 0, \pi_1(v) > 0$ and $C_v^+[a, b] \setminus \{0\} \subseteq (0, \infty) \times \mathbb{R}$. Let l be the integer in the definition of the set A above and define $\hat{U}' := \hat{U} \cup [\hat{U} - (0, l)]$. Suppose for a contradiction that $G^n(\hat{U}) \cap C_v^+(\gamma - \varepsilon, \gamma + \varepsilon) = \emptyset$ for infinitely many $n \in \mathbb{N}$. Slightly reducing ε if necessary, we may assume that $G^n(\hat{U}') \cap C_v^+(\gamma - \varepsilon, \gamma + \varepsilon) = \emptyset$ for infinitely many $n \in \mathbb{N}$. We




Figure 4: Strategy for the proof of Lemma 3.3: The existence of large 'lacunae' in the set A (with blue boundary) forces orbits inside A to deviate far from the line $\mathbb{R}\rho$. In particular, this is true for almost all orbits w.r.t. the measure μ realising the rotation vector $\rho \in \text{Ex}(\rho(G))$. This leads to a contradiction with Lemma 2.3.

first consider the case where $\langle \rho \rangle_v > 0$, such that $\rho \in C_v^+[a, b]$ by Lemma 3.1.

Due to Lemma 2.1 the fact that $\rho_{\hat{U}}(G) \subseteq L_{\lambda, v}$ implies that $\langle G^n z \rangle_v / n$ converges uniformly to λ on $\hat{U}'$ as $n \rightarrow \infty$. Hence, for any $\delta > 0$ there exists $N(\delta) \in \mathbb{N}$ such that

$$(3.6) \quad G^n(\hat{U}') \subseteq S_v[(1 - \delta)n\lambda, (1 + \delta)n\lambda] \quad \forall n \geq N(\delta) .$$

As $\pi_1(v) > 0$, this implies that $\inf \pi_1(G^n(\hat{U}')) \rightarrow \infty$ as $n \rightarrow \infty$. Therefore (3.2) yields that for given $\alpha < a$ and $\beta > b$ and sufficiently large n there holds $G^n(\hat{U}') \subseteq C_v^+[\alpha, \beta]$ (apply (3.2) with $\tilde{\alpha} \in (\alpha, a)$ and $\tilde{\beta} \in (b, \beta)$ to get rid of the constant R). Consequently, if n is large then $G^n(\hat{U}') \cap C_v^+(\gamma - \varepsilon, \gamma + \varepsilon) = \emptyset$ implies $G^n(\hat{U}') \subseteq C_v^+[\alpha, \gamma - \varepsilon]$ or $G^n(\hat{U}') \subseteq C_v^+[\gamma + \varepsilon, \beta]$. We assume that $G^n(\hat{U}') \subseteq C_v^+[\alpha, \gamma - \varepsilon]$ for infinitely many $n \in \mathbb{N}$, the other case is symmetric.

Since $\rho \in \text{Ex}(\rho(G))$, there exists an ergodic measure μ with $\rho_\mu(F) = \rho$ [1, Corollary 3.5]. Let $\Omega \subseteq \mathbb{T}^2$ be such that $g(\Omega) = \Omega$, $\mu(\Omega) = 1$ and

$$(3.7) \quad \lim_{n \rightarrow \infty} (G^n(z) - z)/n = \rho \quad \forall z \in \pi^{-1}(\Omega) .$$

We will show that, in contradiction to Lemma 2.3, for some $s > 0$ there holds

$$(3.8) \quad \forall z \in \Omega \exists n_z \in \mathbb{N} : \quad D_{n_z}^{-\rho^\perp}(z, \rho) \geq s n_z .$$

In order to do so, fix $z \in \Omega$. Let A be as in the proof of Lemma 3.1. Due to (3.7) there exists a lift $z_0 \in A$ of z such that $\pi_1(z_n) \geq K \forall n \in \mathbb{N}$, where $z_n = G^n(z_0)$ and K is chosen as in Claim 3.2. Consequently Claim 3.2 implies that

$$(3.9) \quad z_n \in A \quad \forall n \in \mathbb{N}.$$

Due to (3.6) and the fact that M defined in (3.3) is finite, there exists $\eta > 0$ and $N_1 \in \mathbb{N}$ such that for any $n \geq N_1$ there holds

$$(3.10) \quad G^n(\widehat{U}') \subseteq C_v^+[\alpha, \gamma - \varepsilon] \Rightarrow G^k(\widehat{U}') \subseteq C_v^+[\alpha, \gamma - \varepsilon/2] \quad \forall k \in [(1 - \eta)n, (1 + \eta)n].$$

Now, choose δ in (3.6) sufficiently small, such that $(1 + \delta)(1 - \eta) < 1 < (1 - \delta)(1 + \eta)$. Then choose $N_2 \in \mathbb{N}$ such that for all $n \geq N_2$ there holds

$$(3.11) \quad [\lambda n - M, \lambda n + M] \subseteq [(1 + \delta)(1 - \eta)\lambda n, (1 - \delta)(1 + \eta)\lambda n],$$

$$(3.12) \quad \lambda n - M \geq M \cdot N(\delta) \cdot \|v\| + \sup \langle \widehat{U}' \rangle_v.$$

Suppose $n \geq N_3 := \max\{N(\delta)/(1 - \delta), N_1, N_2\}$ and $G^n(\widehat{U}) \subseteq C_v^+[\alpha, \gamma - \varepsilon]$. Then by combining (3.6) and (3.10)–(3.12) we obtain that

$$(3.13) \quad G^k(\widehat{U}') \cap S_v(\lambda n - M, \lambda n + M) \subseteq C_v^+[\alpha, \gamma - \varepsilon/2] \quad \forall k \in \mathbb{N}.$$

(Treat the cases $k \leq N(\delta)$, $k \in (N(\delta), (1 - \eta)n)$ and $k \geq (1 + \eta)n$ separately to show that for all such k the set $G^k(\widehat{U})$ does not intersect $S_v(\lambda n - M, \lambda n + M)$ and then use (3.10) for the remaining k .) This means in particular that

$$(3.14) \quad A \cap S_v(\lambda n - M, \lambda n + M) \subseteq C_v^+[\alpha, \gamma - \varepsilon/2] \quad \forall n \geq N_3 : G^n(\widehat{U}) \subseteq C_v^+[\alpha, \gamma - \varepsilon].$$

Now, as $\mathbb{R}\rho = C_v[\gamma, \gamma]$ exist constants $r > 0$ and $N_4 \geq N_3$ such that

$$(3.15) \quad \langle z' - z_0 \rangle_{\rho^\perp} \leq -rn \quad \forall n \geq N_4, \quad z' \in S_v(\lambda n - M, \lambda n + M) \cap C_v^+[\alpha, \gamma - \varepsilon/2].$$

Further, since z and its lift z_0 are fixed and due to (3.7) (applied to z_0) there exist constants $N_5 = N_5(z) \geq N_4$ and $c > 0$, with c only depending on ρ, γ and ε , such that

$$(3.16) \quad \forall n \geq N_5 \exists n_z \leq cn : z_{n_z} \in S_v(\lambda n - M, \lambda n + M)$$

(Note that the orbit of z_0 has to pass through $S_v[\lambda n - M, \lambda n + M]$ by definition of M , and due to (3.7) this happens approximately at time $n\lambda/\langle \rho \rangle_v$.) Choose $n \geq N_5$ with $G^n(\widehat{U}) \subseteq C_v^+[\alpha, \gamma - \varepsilon]$. Let n_z be as in (3.16). Since $z_{n_z} \in A$ by (3.9), we obtain $\langle z_{n_z} - z_0 \rangle_{\rho^\perp} \leq -rn$ from (3.14) and (3.15). Thus, if $s = r/c$ then $D_{n_z}^{-\rho^\perp}(z, \rho) \geq sn_z$. Since $z \in \Omega$ was arbitrary, this proves (3.8).

Finally, if $\langle \rho \rangle_v < 0$ then we can proceed in the same way by regarding the inverse of G . Due to the symmetry in the statement of Claim 3.2 we obtain that for a suitable lift z_0 of $z \in \Omega$ the whole backwards orbit of z_0 remains in A , and the remaining argument is exactly the same as before. $\square$

Corollary 3.4. *Suppose $\rho_{\widehat{U}}(G) = L_{\lambda,v}[a, b]$ with $a < b$. Then there exist constants $c > 0$ and $N' \in \mathbb{N}$ such that*

$$(3.17) \quad \sup_{z \in \widehat{U}} \langle G^n(z) \rangle_{v^\perp} - \inf_{z \in \widehat{U}} \langle G^n(z) \rangle_{v^\perp} > cn \quad \forall n \geq N'.$$

Proof. As $L_{\lambda,v}[a, b] = \rho_{\widehat{U}}(G) \subseteq \rho(G)$, the set $\rho(G)$ must have at least two linearly independent extremal points $\rho_1, \rho_2 \neq 0$. Due to Lemma 3.1 these are contained in $C_v[a, b]$, such that $\rho_1 \subseteq C_v[\gamma_1, \gamma_1]$ and $\rho_2 \subseteq C_v[\gamma_2, \gamma_2]$ for some $\gamma_1 \neq \gamma_2 \in \mathbb{R}$. The statement now follows from Lemma 3.3 together with the fact that $\inf \langle G^n(\widehat{U}) \rangle_v \rightarrow \infty$ as $n \rightarrow \infty$. $\square$

We can now describe two situations in which the rotation subset of $\widehat{U}$ determines the whole rotation set completely, or at least forces it to be contained in a line segment.

Lemma 3.5. *Suppose $\rho_{\widehat{U}}(G) = L_{\lambda,v}[a, b]$ with $a < b$. Then $\rho(G) = L_{\lambda,v}[a, b]$.*

Proof. Due to Lemma 3.1, it suffices to show $\rho(G) \subseteq L_{\lambda,v}$. Suppose for a contradiction that $\rho(G) \not\subseteq L_{\lambda,v}$. Then there exists an extremal point $\rho \in \text{Ex}(\rho(G)) \setminus L_{\lambda,v}$. We assume w.l.o.g. that $\|v\| = 1$ and $\langle \rho \rangle_v > \lambda$. Since ρ is realised by an ergodic measure, there exists $z_0 \in \mathbb{R}^2$ with $\lim_{n \rightarrow \infty} (z_n - z_0)/n = \rho$, where $z_n = G^n(z_0)$ as above. Fix $\eta > 0$ and $k_0 \in \mathbb{N}$ such that

$$(3.18) \quad \langle z_k - z_0 \rangle_v > (1 + \eta)\lambda k \quad \forall k \geq k_0.$$

Further, fix $\delta > 0$ such that

$$(3.19) \quad \delta(1 + 15M/c) < \eta,$$

with M defined by (3.3) and c as in Corollary 3.4. Choose $n_0 \in \mathbb{N}$ such that

$$(3.20) \quad G^n(\widehat{U}) \subseteq S_v[(1 - \delta)\lambda n, (1 + \delta)\lambda n] \quad \forall n \geq n_0 \quad \text{and}$$

$$(3.21) \quad \delta\lambda n_0 \geq 2.$$

Then choose $k \geq k_0$ and $n \geq n_0$ such that

$$(3.22) \quad (4M + 2)k \leq cn \leq 5Mk.$$




Figure 5: Proof of Lemma 3.5: The slow movement of the curve Γ impedes a faster movement of the point z_0 under iteration.

Due to Corollary 3.4 there exists a simple arc $\Gamma_0 \subseteq G^n(\widehat{U})$ with endpoints $\zeta_1, \zeta_2 \in G^n(\widehat{U})$ such that $\langle \zeta_2 - \zeta_1 \rangle_{v^\perp} > cn$ and $\Gamma_0 \subseteq S_{v^\perp}[\langle \zeta_1 \rangle_{v^\perp}, \langle \zeta_2 \rangle_{v^\perp}]$. Let Γ_1 and Γ_2 be properly embedded half-lines (proper images of $\mathbb{R}^+$) which join ζ_1 , respectively ζ_2 , to infinity and satisfy

$$(3.23) \quad \Gamma_i \setminus \{z_i\} \cap S_{v^\perp}[\langle \zeta_1 \rangle_{v^\perp}, \langle \zeta_2 \rangle_{v^\perp}] = \emptyset.$$

Let the properly embedded line $\Gamma = \Gamma_0 \cup \Gamma_1 \cup \Gamma_2$ be oriented in the direction from ζ_1 to ζ_2 and denote by W the connected component of $\mathbb{R}^2 \setminus \Gamma$ to the left of Γ . Then, since $\Gamma_0 \subseteq G^n(\widehat{U})$ by assumption and due to (3.20) and (3.23), W contains the set $S_v(-\infty, (1 - \delta)\lambda n] \cap S_{v^\perp}[\langle \zeta_1 \rangle_{v^\perp}, \langle \zeta_2 \rangle_{v^\perp}]$. Further, due to (3.21) and (3.22) the set

$$(3.24) \quad W_0 = S_v[(1 - 2\delta)\lambda n, (1 - \delta)\lambda n] \cap S_{v^\perp}[\langle \zeta_1 \rangle_{v^\perp} + 2kM, \langle \zeta_2 \rangle_{v^\perp} - 2kM]$$

is a rectangle whose side-lengths are greater than 2. Hence W_0 contains a fundamental domain of $\mathbb{T}^2$ and by replacing z_0 with an integer translate if necessary we may assume $z_0 \in W_0$. This implies in particular that $\langle z_0 \rangle_v \geq (1 - 2\delta)\lambda n$. Now, there holds

$$(3.25) \quad z_k \in G^k(W) \cap S_{v^\perp}[\langle \zeta_1 \rangle_{v^\perp} + kM, \langle \zeta_2 \rangle_{v^\perp} - kM].$$

However, due to (3.20) and (3.23) we have

$$(3.26) \quad G^k(W) \cap S_v((1+\delta)\lambda(n+k), \infty) \cap S_{v^\perp}[\langle \zeta_1 \rangle_{v^\perp} + kM, \langle \zeta_2 \rangle_{v^\perp} - kM] = \emptyset,$$

such that $z_k \in S_v(-\infty, (1+\delta)\lambda(n+k)]$. Thus, using (3.22) and (3.19) we obtain $\langle z_k - z_0 \rangle_v \leq (1+\delta)\lambda(n+k) - (1-2\delta)\lambda n < (1+\eta)\lambda k$, in contradiction to (3.18). $\square$

More or less along the same lines we obtain the following.

Lemma 3.6. *Suppose $\rho_{\widehat{U}}(G) \subseteq \mathbb{R} \cdot v$ is a line segment of positive length. Then $\rho(G) \subseteq \mathbb{R} \cdot v$.*

Proof. Suppose for a contradiction that $\rho(G) \not\subseteq \mathbb{R} \cdot v$ and assume without loss of generality that there exists $\rho \in \text{Ex}(\rho(G))$ with $\sigma := \langle \rho \rangle_{v^\perp} > 0$. Then as above, there exists a point $z_0 \in \mathbb{R}^2$ with $\lim_{n \rightarrow \infty} (z_n - z_0)/n = \rho$, where $z_n = G^n(z_0)$. Hence, there exists an integer $N \geq 1$ such that

$$(3.27) \quad \langle z_n - z_0 \rangle_{v^\perp} > n\sigma/2 \quad \forall n \geq N.$$

Let av and bv be the endpoints of $\rho_U(G)$, with $a < b$. Then by definition of $\rho_{\widehat{U}}(G)$ in (1.2) and due to the connectedness of $\bigcup_{j=0}^m G^j(\widehat{U})$, there are infinitely many $m \in \mathbb{N}$ such that there exists a simple arc $\Gamma_0 \subseteq \bigcup_{j=0}^m G^j(\widehat{U})$ with endpoints ζ_1 and ζ_2 such that

$$(3.28) \quad \langle \zeta_2 - \zeta_1 \rangle_v \geq m(b-a)/3.$$

Of course, we may assume $\Gamma_0 \subseteq S_v[\langle \zeta_1 \rangle_v, \langle \zeta_2 \rangle_v]$. Further, given any $\delta > 0$ there exists $m_0 \in \mathbb{N}$ such that for all $m \geq m_0$ we have

$$(3.29) \quad G^n(\Gamma_0) \subseteq \bigcup_{j=0}^{n+m} G^j(\widehat{U}) \subseteq S_{v^\perp}[-\delta(m+n), \delta(m+n)] \quad \forall n \in \mathbb{N}.$$

Now, fix $\delta > 0$ such that

$$(3.30) \quad 4\delta/(\sigma - 2\delta) < (b-a)/12$$

and choose $m \geq m_0$ such that there exists an integer $n_0 \geq N$ with

$$(3.31) \quad n_0 \in \left[\frac{4\delta m + 4}{\sigma - 2\delta}, \frac{m(b-a) - 6}{12M} \right].$$

Let Γ_1 and Γ_2 be properly embedded half-lines such that Γ_i joins ζ_i to infinity and $\Gamma_i \setminus \{\zeta_i\}$ is disjoint from $S_v[\langle \zeta_1 \rangle_v, \langle \zeta_2 \rangle_v]$. Let $\Gamma = \Gamma_0 \cup \Gamma_1 \cup \Gamma_2$ be oriented in the direction from ζ_1 to ζ_2 . Denote by W the connected component of $\mathbb{R}^2 \setminus \Gamma$ to the right of Γ . Then $W_0 = S_v[\langle \zeta_1 \rangle_v + 2n_0M, \langle \zeta_2 \rangle_v - 2n_0M] \cap S_{v^\perp}[-\delta m - 2, -\delta m] \subseteq W$ contains a fundamental domains of $\mathbb{T}^2$. (Note that due to (3.28) and (3.31) there holds $\langle \zeta_2 \rangle_v - \langle \zeta_1 \rangle_v \geq 4n_0M + 2$.) Replacing z_0 by an integer translate if necessary, we may therefore assume $z_0 \in W_0$. It follows that

$$(3.32) \quad z_{n_0} \in G^{n_0}(W) \cap S_v[\langle \zeta_1 \rangle_v + n_0M, \langle \zeta_2 \rangle_v - n_0M].$$

However, due to (3.29) and the choice of Γ the set $G^{n_0}(W)$ is disjoint from $S_v[\langle \zeta_1 \rangle_v + n_0M, \langle \zeta_2 \rangle_v - n_0M] \cap S_{v^\perp}(\delta(m+n_0), \infty)$, such that

$$(3.33) \quad \langle z_{n_0} - z_0 \rangle_{v^\perp} \leq \delta(2m + n_0) + 2.$$

However, from (3.31) we obtain that $\delta(2m + n_0) + 2 \leq n_0\sigma/2$, such that (3.33) contradicts (3.27). $\square$

4 Proof of the main results

The following basic observation will be used in the proof of Theorem 1.1. Recall that $\varphi_n(z) = (F^n(z) - z)/n$.

Lemma 4.1. *Suppose F is a lift of $f \in \text{Homeo}_0(\mathbb{T}^d)$ and $U \subseteq \mathbb{T}^d$ is open, connected, bounded and recurrent. Further, assume that $\rho_U(F) = \mathcal{S} \subseteq \mathbb{R} \cdot v$ is a line segment with $0 \notin \mathcal{S}$ and v, v' are linearly independent. Let $\widehat{U}$ be a connected component of $\pi^{-1}(U)$. Then there exists $p \in \mathbb{N}$ and $w \in \mathbb{Z}^2$ linearly independent of v' such that*

$$(4.1) \quad (F^p(\widehat{U}) - w) \cap \widehat{U} \neq \emptyset.$$

In particular, if v is not the scalar multiple of an integer vector then there exist infinitely many pairs (p_i, w_i) with pairwise independent integer vectors w_i that satisfy (4.1).

Proof. Choose $\varepsilon > 0$ such that $B_{2\varepsilon}(\rho_U(F)) = B_{2\varepsilon}(\mathcal{S})$ is disjoint from $\mathbb{R}v'$. As $\widehat{U}$ is bounded, there exists $n_0 \in \mathbb{N}$ such that

$$\frac{1}{n} \left(F^n(\widehat{U}) - \widehat{U} \right) \subseteq B_\varepsilon(\varphi_n(\widehat{U})) \quad \forall n \geq n_0.$$

(Here $A - B = \{z - z' \mid z \in A, z' \in B\}$.) Due to Lemma 2.1 we may further assume, by increasing n_0 if necessary, that

$$\varphi_n(\widehat{U}) \subseteq B_\varepsilon(\rho_U(F)) \quad \forall n \geq n_0.$$

We thus obtain

$$\left(F^n(\widehat{U}) - \widehat{U} \right) \cap \mathbb{R}v' = \emptyset \quad \forall n \geq n_0.$$

Now, since U is recurrent, there exists $p \geq n_0$ with $f^p(U) \cap U \neq \emptyset$ and hence an integer vector w with $(F^p(\widehat{U}) - w) \cap \widehat{U} \neq \emptyset$. As w belongs to $F^n(\widehat{U}) - \widehat{U}$ it must be linearly independent of v' . $\square$

Proof of Theorem 1.1. We start with (b) and (c) and prove (a) at the end.

(b) Let f, F, U and $\mathcal{S} = \rho_U(F)$ be as in the statement of the theorem. Let $\widehat{U}$ be a connected component of $\pi^{-1}(U)$. We first assume that the line passing through $\mathcal{S}$ does not contain any rational points. Since U is non-wandering, there exists $p \in \mathbb{N}$ and $w \in \mathbb{Z}^2$ such that $(F^p(\widehat{U}) - w) \cap \widehat{U} \neq \emptyset$. Let $G = F^p - w$. Then $\rho_{\widehat{U}}(G) = p\mathcal{S} - w = L_{\lambda, v}[a, b]$ for some $\lambda \in \mathbb{R}$, $v \in \mathbb{R}_*^2$ and $a < b$. Further, the line $L_{\lambda, v}$ contains no rational points either and therefore $\lambda \neq 0$. Hence $g = f^p$, G and $\widehat{U}$ satisfy the assumptions of Lemma 3.5 and we obtain $\rho(G) = L_{\lambda, v}[a, b]$ and thus $\rho(F) = (\rho(G) + w)/p = \mathcal{S}$. It remains to treat the case where the line passing through $\mathcal{S}$ has irrational slope and contains a single rational point. We choose $\widehat{U}$ and G as above and again have $\rho_{\widehat{U}}(G) = L_{\lambda, v}[a, b]$ for some $\lambda \in \mathbb{R}$, $v \in \mathbb{R}_*^2$ and $a < b$. If $\lambda \neq 0$, then we can proceed exactly as before. However, this time we may have $\lambda = 0$ since $L_{\lambda, v}$ may pass through 0. In this case Lemma 4.1 yields the existence of a pair $\tilde{p} \in \mathbb{N}$ and $\tilde{w} \in \mathbb{Z}^2$ such that $\tilde{w}$ and w are linearly independent and $\tilde{G}(\widehat{U}) \cap \widehat{U} \neq \emptyset$, where $\tilde{G} = F^{\tilde{p}} - \tilde{w}$. Then

$$\rho_{\widehat{U}}(\tilde{G}) = \tilde{p} \cdot \rho_{\widehat{U}}(F) - \tilde{w} = \tilde{p} \cdot \left(\frac{\rho_{\widehat{U}}(G) + w}{p} \right) - \tilde{w} \subseteq L_{0, v} + \left(\frac{\tilde{p}}{p} \cdot w - \tilde{w} \right)$$

As w and $\tilde{w}$ are linearly independent there holds $\frac{\tilde{p}}{p} \cdot w - \tilde{w} \neq 0$. At the same time, this vector is not in $L_{0, v} = \{v\}^\perp$ since the only rational vector contained in this line is 0. Therefore $\rho_{\widehat{U}}(\tilde{G}) = L_{\tilde{\lambda}, v}$ for some $\tilde{\lambda} \neq 0$ and we can apply Lemma 3.5 to $\tilde{g} = f^{\tilde{p}}, \tilde{G}$ and $\widehat{U}$ to obtain $\rho_{\widehat{U}}(F) = \mathcal{S}$ as before.

(c) Suppose $\rho_U(F) = \{\rho\}$ with ρ irrational. As above, we choose $p \in \mathbb{N}$, $w \in \mathbb{Z}^2$ and $G = F^p - w$ such that $G(\widehat{U}) \cap \widehat{U} \neq \emptyset$. Then $\rho_{\widehat{U}}(G) = \{p\rho - w\} = L_{1, p\rho - w}[0, 0]$. By Lemma 3.1 all extremal points of $\rho(G)$ are contained in $C_{p\rho - w}[0, 0] = \mathbb{R} \cdot (p\rho - w)$ and hence $\rho(G) \subseteq \mathbb{R} \cdot (p\rho - w)$. This implies $\rho(F) \subseteq \mathbb{R} \cdot (\rho - w/p) + w/p =: A_1$.

Due to Lemma (4.1) we can repeat this argument with a second pair $\tilde{p} \in \mathbb{N}$ and $\tilde{w} \in \mathbb{Z}^2$, with $\tilde{w}$ linearly independent of w , and obtain $\rho(F) \subseteq \mathbb{R} \cdot (\rho - \tilde{w}/\tilde{p}) + \tilde{w}/\tilde{p} =: A_2$. It follows that $\rho(F)$ is contained in the intersection of the two lines A_1 and A_2 , which is equal to $\{\rho\}$. (Note that as ρ is irrational and the vectors w/p and $\tilde{w}/\tilde{p}$ are both rational and linearly independent, the vectors $\rho - w/p$ and $\rho - \tilde{w}/\tilde{p}$ are linearly independent as well.)

(a) Due to (a) and (b), it only remains to treat the two cases where $\rho_U(F)$ is either a line segment of positive length containing a rational vector or $\rho_U(F)$ is reduced to a single semi-rational vector. The second case is treated exactly as in (b), the only difference is that we cannot necessarily repeat the argument with a second $\tilde{w}$ to conclude that $\rho(F)$ is a singleton.

Hence, suppose that $\rho_U(F) \subseteq \mathbb{R} \cdot v$ is a line segment of positive length and $\rho_U(F)$ contains a rational. By going over to a suitable iterate, we may assume w.l.o.g. that $0 \in \rho_U(F)$. Again, there exists a pair $p \in \mathbb{N}$ and $w \in \mathbb{Z}^2$ such that $G(\widehat{U}) \cap \widehat{U} \neq \emptyset$, where $G = F^p - w$. We have $\rho_{\widehat{U}}(G) \subseteq L_{\lambda, v^\perp}$ with $\lambda := \langle w \rangle_{v^\perp}$. If $\lambda \neq 0$, then Lemma 3.5 implies that $\rho(G) = \rho_{\widehat{U}}(G)$ and hence $\rho(F) = \rho_U(F)$. If $\lambda = 0$, such that $\rho_{\widehat{U}}(G) \subseteq \mathbb{R} \cdot v$, then Lemma 3.6 implies $\rho(G) \subseteq \mathbb{R} \cdot v$, such that we obtain $\rho(F) = \rho(G)/p \subseteq \mathbb{R} \cdot v$ as well. $\square$

Proof of Theorem 1.3. Suppose $U \subseteq \mathbb{T}^2$ is open, bounded and connected. As f is non-wandering, U is also recurrent. Suppose $\text{Conv}(\rho_U(F))$ has empty interior, that is, $\rho_U(F)$ is contained in a line. Then Theorem 1.1 implies that $\rho_U(F)$ is reduced to a single rational vector. By going over to a suitable iterate, we may assume $\rho_U(F) = \{0\}$. Let $\mathcal{D} := \{z \in \mathbb{T}^2 \mid \exists \varepsilon > 0 : \rho_{B_\varepsilon(z)}(F) = 0\}$ and note

that $U \subseteq \mathcal{D}$. Let D denote the connected component of $\mathcal{D}$ that contains U . Since $\mathcal{D}$ is f -invariant and f is non-wandering, D is periodic with period p for some $p \in \mathbb{N}$.

We claim that D contains no essential simple closed curve. In order to see this, suppose for a contradiction that the curve $\gamma \subseteq D$ is essential with homotopy vector $v \in \mathbb{Z}_*^2$. Let Γ be a connected component of $\pi^{-1}(\gamma)$. Then Γ is a properly embedded line that remains in a bounded distance of $\mathbb{R} \cdot v$. Furthermore, due to compactness $\gamma \subseteq D$ is covered by a finite number of open sets U_i with $\rho_{U_i}(F) = \{0\}$, which implies $\rho_\gamma(F) = \{0\}$. However, this clearly contradicts the existence of rotation vectors ρ with $\langle \rho \rangle_v \neq 0$ in $\rho(F)$. (Compare, for example, the proof of Lemma 3.6.)

In a similar way, we see that D is simply-connected. If $\Gamma \subseteq \pi^{-1}(D)$ is a closed Jordan curve, then by compactness we obtain $\rho_\Gamma(F) = \{0\}$. Consequently the Jordan domain $J(\Gamma)$ bounded by Γ has rotation subset $\rho_{J(\Gamma)}(F) = \{0\}$ and therefore belongs to $\pi^{-1}(D)$ as well. Thus D is the required f^p -invariant topological disk. Finally, as D is homeomorphic to $\mathbb{R}^2$, the restriction $f|_D^p$ defines a plane homeomorphism. Since $f|_D^p$ is non-wandering, it must have a fixed point. (This follows, for example, from the Brouwer Plane Translation Theorem or from Franks Lemma [11]). $\square$

Proof of Proposition 1.5.

(a) Suppose $f \in \text{Homeo}_0(\mathbb{T}^2)$ is non-wandering, F is a lift and $z \in \mathcal{C}(f)$ is $\frac{1}{2}$ -Lyapunov stable. Choose $\delta > 0$ such that $f^n(B_\delta(z)) \subseteq B_{\frac{1}{2}}(f^n(z)) \forall n \in \mathbb{N}$. Let $z_0 \in \mathbb{R}^2$ be a lift of z and $\widehat{U} := B_\delta(z_0)$. Then $F^n(\widehat{U}) \subseteq B_{\frac{1}{2}}(F^n(z_0)) \forall n \in \mathbb{N}$, in particular $\text{diam}(F^n(\widehat{U})) \leq 1$.

Since f is non-wandering U is recurrent, such that there exist infinitely many pairs $(p, w) \in \mathbb{N} \times \mathbb{Z}^2$ with $(F^p(\widehat{U}) - w) \cap \widehat{U} \neq \emptyset$. The latter implies $F^p(\widehat{U}) \subseteq B_{1+\delta}(z_0 + w)$. Further, as $(F^{np}(\widehat{U}) - w) \cap F^{(n-1)p}(\widehat{U}) \neq \emptyset \forall n \in \mathbb{N}$, we obtain by induction that

$$F^{np}(\widehat{U}) \subseteq B_{n+\delta}(z_0 + nw) \quad \forall n \in \mathbb{N}.$$

However, this implies $\rho_{\widehat{U}}(F) \subseteq B_{\frac{1}{p}}(w)$, such that $\text{diam}(\rho_{\widehat{U}}(F)) \leq 1/p$. As p can be chosen arbitrarily large we obtain $\text{diam}(\rho_{\widehat{U}}(F)) = 0$, in contradiction to $z \in \mathcal{C}(f)$.

(b) Suppose f is area-preserving, U is a connected and bounded neighbourhood of $z \in \mathcal{C}(f)$ and $\widehat{U}$ is a connected component of $\pi^{-1}(U)$. Birkhoffs Ergodic Theorem implies that Lebesgue-almost every point $z' \in \widehat{U}$ has a rotation vector $\rho(F, z') = \lim_{n \rightarrow \infty} (F^n(z') - z')/n$ (that is, the limit exists). If $\lim_{n \rightarrow \infty} \text{diam}(\langle F^n(\widehat{U}) \rangle_v)/n = 0$ this implies $\rho_{\widehat{U}}(F) \subseteq L_{\lambda, v}$ with $\lambda = \langle \rho(F, z') \rangle_v$, contradicting $z \in \mathcal{C}(f)$. $\square$

5 A parameter family of Misiurewicz-Ziemian type

Examples of a toral homeomorphisms, homotopic to the identity, whose rotation set has non-empty interior were introduced by Misiurewicz and Ziemian [3] via lifts of the form

$$(5.1) \quad F(x, y) = (x + \psi_2(y + \psi_1(x)), y + \psi_1(x)) ,$$

with continuous and 1-periodic functions $\psi_i : \mathbb{R} \rightarrow \mathbb{R}$, $i = 1, 2$. When $\psi_1(0) = \psi_2(0) = 0$ and $\psi_1(\frac{1}{2}) = \psi_2(\frac{1}{2}) = 1$, it can be easily checked that the points $(0, 0)$, $(\frac{1}{2}, 0)$, $(0, \frac{1}{2})$ and $(\frac{1}{2}, \frac{1}{2})$ are fixed under the induced map $f \in \text{Homeo}_0(\mathbb{T}^2)$ and have rotation vectors $(0, 0)$, $(0, 1)$, $(1, 0)$ and $(1, 1)$, respectively. Since the rotation set $\rho(F)$ is convex, it contains the square $[0, 1]^2$. More or less the same type of examples was proposed independently by Llibre and MacKay in [5].

In order to give a smooth example, we slightly modify this structure and let $\psi_1(x) = \alpha \sin(2\pi x)$ and $\psi_2(y) = \beta \sin(2\pi y)$ to obtain the parameter family

$$(5.2) \quad F_{\alpha, \beta}(x, y) = (x + \beta \sin(2\pi(y + \alpha \sin(2\pi x))), y + \alpha \sin(2\pi x)) .$$

We denote the toral diffeomorphisms induced by these lifts by $f_{\alpha, \beta}$. We note that $f_{\alpha, \beta}$ is area-preserving for all $\alpha, \beta \in \mathbb{R}$ and hence, in particular, non-wandering.

For $F_* = F_{\frac{1}{2}, \frac{1}{2}}$, the points $(\frac{1}{4}, 0)$, $(0, \frac{1}{4})$, $(\frac{3}{4}, 0)$ and $(0, \frac{3}{4})$ are 2-periodic with rotation vectors $(0, \frac{1}{2})$, $(\frac{1}{2}, 0)$, $(0, -\frac{1}{2})$ and $(-\frac{1}{2}, 0)$. The rotation set $\rho(F_*)$ therefore contains the square $Q = \{(x, y) \mid |x| + |y| \leq \frac{1}{2}\}$ spanned by these vectors. Hence, the toral diffeomorphism $f_* = f_{\frac{1}{2}, \frac{1}{2}}$ induced by F_* satisfies all the assumptions of Theorem 1.1. Furthermore, this remains true for all (α, β) in a neighbourhood of $(\frac{1}{2}, \frac{1}{2})$. The reason is that when the rotation set has non-empty interior, then it depends continuously on the toral homeomorphism in the C^0 -topology [3].

The points $(\frac{i}{4}, \frac{j}{4})$ with $i, j = 1, 3$ are 2-periodic for f_* and have rotation vector $(0, 0)$. The phase portrait of f_* in Figure 1⁴ clearly suggests that these points are surrounded by (star-shaped) elliptic islands. Since the differential DF_*^2 in these points is the identity matrix, it is difficult to establish the existence of elliptic islands in a rigorous way by the application of standard KAM-results⁵, and we shall not further pursue this issue here. Yet, the pictures obtained by simulations show typical features of a KAM-type elliptic region, with invariant curves prevalent in the centre and at least one clearly visible instability zone towards the boundary (Figure 6).




Figure 6: Enlargement of the elliptic island around $(\frac{1}{4}, \frac{1}{4})$ in Figure 1. Left: 20000 iterates of the starting points $(0.253 + i \cdot 0.00455) \cdot (1, 1)$, $i = 1, \dots, 10$. Right: 100000 iterates of the starting point $0.298429 \cdot (1, 1)$ in the instability region.)

Instead, we want to close by briefly discussing some of the symmetries present in Figure 1. The picture is invariant under the rotation with angle $\pi/2$ around the point $(\frac{1}{2}, \frac{1}{2})$, which is given by the map $R(x, y) = (-y, x)$ on the torus. However, f_* is not conjugate to itself by R . The reason why this symmetry nevertheless appears is the fact that R conjugates f_* to its inverse, that is, $R^{-1} \circ f_* \circ R = f_*^{-1}$, and for the visualisation of elliptic islands it does not make any difference if f_* is replaced by f_*^{-1} . This remains true for $f_{\alpha, \alpha}$ for all α , since

$$(5.3) \quad R^{-1} \circ f_{\alpha, \alpha} \circ R = f_{\alpha, \alpha}^{-1} \quad \forall \alpha \in \mathbb{R}.$$

Two symmetries conjugating $f_{\alpha, \beta}$ to itself, for all $\alpha, \beta \in \mathbb{R}$, are the rotation $S(x, y) = (-x, -y)$ with angle π and the map $T(x, y) = (x + \frac{1}{2}, -y + \frac{1}{2})$, which is the reflexion along the x -axis composed with the shift by $(\frac{1}{2}, \frac{1}{2})$. This implies that if the points $(\frac{i}{4}, \frac{j}{4})$ with $i, j = 1, 3$ are surrounded by elliptic islands, then these are all isometric to each other, but the isometries are orientation-reversing if (i, j) and (i', j') are such that $i = i'$ or $j = j'$. When $\alpha = \beta$ the additional symmetry given by R implies that the elliptic islands are self-symmetric. For the island surrounding $(\frac{1}{4}, \frac{1}{4})$ the symmetry axis is $L = \{(x, y) \mid x + y = \frac{1}{2}\}$, for the others it is the respective image of L under R^i , $i = 1, 2, 3$.

When $\alpha \neq \beta$ this self-symmetry of the elliptic islands breaks down since (5.3) does not hold anymore. With a slight adjustment of the parameters and some imagination in the Rorschach test below, this allows to return to a more aquatic environment (Figure 7).

References

- [1] M. Misiurewicz and K. Ziemian. Rotation sets for maps of tori. *J. Lond. Math. Soc.*, 40:490–506, 1989.
- [2] J. Franks. Realizing rotation vectors for torus homeomorphisms. *Trans. Am. Math. Soc.*, 311(1):107–115, 1989.

⁴Picture produced with SCILAB by computing 1000 iterates for each starting point in a 40x40-grid on T^2 and omitting the first 100 iterates for the plot.

⁵This would require an elliptic differential matrix with irrational rotation angle [9].




Figure 7: Elliptic sharks circling a chaotic ocean. (Phase portrait of $f_{\alpha,\beta}$ with $\alpha = 0.5$ and $\beta = 0.502$, with the same indications as in Footnote 4.)

- [3] M. Misiurewicz and K. Ziemian. Rotation sets and ergodic measures for torus homeomorphisms. *Fundam. Math.*, 137(1):45–52, 1991.
- [4] T. Jäger. Linearisation of conservative toral homeomorphisms. *Invent. Math.*, 176(3):601–616, 2009.
- [5] J. Llibre and R.S. MacKay. Rotation vectors and entropy for homeomorphisms of the torus isotopic to the identity. *Ergodic Theory Dyn. Syst.*, 11:115–128, 1991.
- [6] J. Moser. On invariant curves of area-preserving mappings of an annulus. *Nachr. Akad. Wiss. Göttingen, II Math. Phys. Kl.*, pages 1–20, 1962.
- [7] M. Herman. Sur les courbes invariantes par les difféomorphismes de l’anneau. *Asterisque*, 103–104, 1983.
- [8] H. Rüssmann. On the existence of invariant curves of twist mappings of the annulus. In *Geometric dynamics*, Proc. Int. Symp. Rio de Janeiro/Brasil 1981. *Lect. Notes Math.* 1007, pages 677–718, 1983.
- [9] A. Katok and B. Hasselblatt. *Introduction to the Modern Theory of Dynamical Systems*. Cambridge University Press, 1997.
- [10] J. Kwapisz. Every convex polygon with rational vertices is a rotation set. *Ergodic Theory Dyn. Syst.*, 12:333–339, 1992.
- [11] J. Franks. Generalizations of the Poincaré-Birkhoff theorem. *Ann. Math. (2)*, 128(1):139–151, 1988.